

Spécialité de Master « Optique, Matière, Paris »

Proposition de stage 2016-2017

Laboratoire: Laboratoire Aimé Cotton

Adresse: Campus d'Orsay, Bâtiment 505, Rue du Belvédère, 91 405 Orsay

Directeur du laboratoire: J.-F. Roch

<http://www.lac.u-psud.fr>



Responsable(s) du stage: E. Deleporte et D. Garrot

Téléphone: E. Deleporte: 01 69 35 20 24, D. Garrot: 01 39 25 46 03


e-mail: Emmanuelle.Deleporte@ens-paris-saclay.fr, Damien.Garrot@uvsq.fr

Dielectric and quantum confinement in mixed-dimensionality 2D/3D hybrid organic-inorganic perovskites

Scientific project:

Since 2012, the 3D hybrid organic-inorganic (3D HOP) perovskites $\text{CH}_3\text{NH}_3\text{PbX}_3$ (X: I, Br, Cl) represents a “material breakthrough” for photovoltaics: in only 3 years, the efficiency of $\text{CH}_3\text{NH}_3\text{PbI}_3$ based solar cells has progressed from 12% to 22,1%.⁽¹⁾ However, one of the main technological issue to overcome is the stability of the material, especially its sensitivity to moisture. The solution might come from the layered 2D HOP and the mixed 2D/3D structures. **In particular, photovoltaic devices based on the mixed 2D/3D HOP have demonstrated superior stability and a growing efficiency.**⁽²⁾

The 2D/3D HOP family has the following chemical formula $(\text{RNH}_3)_2(\text{CH}_3\text{NH}_3)_{m-1}\text{Pb}_m\text{X}_{3m+1}$ (R: organic group, X: halogen). The variable m corresponds to the number of inorganic layers intercalated between the organic molecules (see figure). For $m=1$, the material adopts a two-dimensional structure (2D HOP), where carriers are confined in an atomically thin inorganic layer, made of PbX_6 octahedras, and form strongly bound excitons (exciton binding energy of several 100 meV).⁽³⁾ At the limit case of $m=\infty$, we obtain the 3D hybrid perovskite structure $\text{CH}_3\text{NH}_3\text{MX}_3$ for which the exciton binding energy is only of a few meV and the transport properties are dominated by free carriers at room temperature.⁽⁴⁾ **From $m=1$ (2D) to $m=\infty$ (3D), the dimensionality and thus the excitonic properties of HOPs can be tailored.**



Nevertheless the charge carrier properties of these HOP are largely unknown for the moment. Contrary to standard quantum wells, the high contrast between the dielectric constants of the organic and inorganic part generates a dielectric confinement of the carriers, making the physics of the excitons more complex.^(5–7) Besides, Coulombic interaction between the carriers and the dipole moment of the organic cation CH_3NH_2^+ lead to the formation of specific polaronic states.⁽⁸⁾

1. http://www.nrel.gov/ncpv/images/efficiency_chart.jpg

3. Gauthron K. *et al. Opt Express.* **2010**;18(6),5912–9.

5. Abdel-Baki K *et al. J. App. Phys.* **2016**. p. 64301

7. Yangui A, *et al. J. Phys. Chem. C;* **2015**, 119(41), 23638

2. Tsai H *et al. Nature.* **2016**, 536, 312-316.

4. Miyata A. *et al. Nature Physics* **2015**, 11(7), 582

6. Lanty G *et al. J. Phys. Chem. Lett.* **2014**, 5(22),3958.

8. Zhu H *et al. Science.* **2016**, 353,6306

Methods and techniques: Excitonic properties of 2D and 2D/3D HOP single crystals will be studied thanks to linear and non-linear optical experiments. Absorption, photoluminescence (PL) and PL excitation experiments as function of the temperature will give access to exciton binding energies, excitonic structure and electron-phonon coupling. The relationship between the structure, the phase transition and the optical properties will be investigated. Time-resolved spectroscopy will be used to investigate the dynamics of the carrier relaxation.

Possibility to go on with a PhD ? Yes

Envisaged fellowship ? Applying to Ecole Doctorale financial support